LEPTOTYPHLOPS AND RAMPHOTYPHLOPS. FITZINGER, 1843 (REPTILIA, SERPENTES): PROPOSED CONSERVATION UNDER THE PLENARY POWERS. Z.N.(S.) 2155.

By Andrew F. Stimson (*British Museum* (*Natural History*), London, SW7); Joan Robb (*Department of Zoology, University of Auckland, New Zealand*) and Garth Underwood (*Department of Biological Sciences, City of London Polytechnic*)

The present note deals with the names of two genera of burrowing snakes, one in the family TYPHLOPIDAE, the other in the family LEPTOTYPHLOPIDAE.

- 2. Robb (1966) split the genus *Typhlops* Oppel (1811: 54) on the basis of the structure of the male reproductive organs. Those species with hollow, eversible male organs were left in the genus *Typhlops*, while species with solid, protrusible male organs and blind pouches opening from the wall of the cloaca were placed in a separate genus. Robb selected *Ramphotyphlops* Fitzinger (1843: 24, type-species by original designation *Typhlops multilineatus* Schlegel, 1839: 40), as being the oldest available nominal genus whose type-species qualified for inclusion in the newly defined genus.
- 3. McDowell (1974: 20) accepted Robb's splitting of *Typhlops* into two genera, but rejected the name *Ramphotyphlops* on the grounds that it is antedated by *Typhlina* Wagler (1830: 196). McDowell believed that the type species of *Typhlina* was *Acontias lineatus* Schlegel (1839: 39) by subsequent designation (Fitzinger 1843: 24).
- 4. The genus *Typhlina*, when first erected, was based on two species, viz:

Acontias lineatus Reinw.

Typhlops sentemstriatus Schneid.

Acontias lineatus was at the time a nomen nudum. It had previously been published only by Fitzinger (1826: 54), Schlegel (1827: 291), and Boie (1827: 563) and these authors, like Wagler, had given no valid indication as defined in Article 16 of the Code. Thus it was an unavailable name in the sense of Article 68(c) and its designation as type-species of Typhlina is invalid. The only other originally included species, "Typhlops sentemstriatus Schneid". (a lapsus or an unjustified emendation of Anguis septemstriatus Schneider, 1801: 341) had at the time an available name and is therefore the type of Typhlina by monotypy.

5. Thus McDowell's claim that *Typhlina* antedates *Ramphotyphlops* cannot be substantiated, as the type-species of *Typhlina* (septemstriatus) does not come within the ambit of the genus *Ramphotyphlops*.

- 6. This raises another problem. Anguis septemstriatus has long been considered a member of the genus Leptotyphlops Fitzinger (1843: 24, type species by original designation Typhlops nigricans Schlegel, 1839: 38). Typhlina is therefore a senior subjective synonym of Leptotyphlops and would replace that name if the Law of Priority were applied.
- 7. After its original description Leptotyphlops was unused as a valid name for 48 years, being universally treated as a junior synonym of Stenostoma Wagler (in Spix 1824: 68; non Stenostoma Latreille, 1810: 217). When Stenostoma was found to be preoccupied (Boulenger, 1890: 243), there followed some 40 years of instability. Steineger (1891: 501) resurrected Leptotyphlops while Boulenger (1892: 10), in the belief that Leptotyphlops was a nomen nudum, preferred to use Glauconia Gray (1845:139). For three decades Glauconia was more widely used than Leptotyphlops, but during the late twenties the use of the latter name became increasingly popular, so that by 1930 very few specialists were still using Glauconia. Since 1930 Glauconia has appeared in print no more than half a dozen times, while Leptotyphlops has been employed in well over 300 publications including the well known faunal lists and textbooks of Ditmars (1939: 171), Smith & Taylor (1945: 20), Parker (1949: 19), Schmidt (1953: 154), Witte (1953: 152), Loveridge (1957: 245), Wright & Wright (1957: 36), FitzSimons (1962: 77), Roze (1966: 39), Peters & Orejas Miranda (1970: 165), Goin & Goin (1971: 301), Mertens (1971: 76), Leviton (1971: 149), Porter (1972: 211) and Pitman (1974: 65). After 50 years of stability it is felt that no useful purpose would be served by replacing Leptotyphlops with Typhlina, a name unused in its correct sense since its original description.
- 8. This undesirable nomenclatural change could be prevented by suppressing *Typhlina* and placing it on the Official Index of Rejected and Invalid Generic Names in Zoology. However, it could be argued, since the synonymy of this name with *Leptotyphlops* is subjective, that to suppress *Typhlina* would create problems should some future worker decide that *Anguis septemstriatus* and *Typhlops nigricans* are not congeneric. Fortunately *Saigonodon* (Peters, 1881: 71, type-species by original designation *Anguis septemstriatus* Schneider), a junior objective synonym of *Typhlina*, is available and could be utilised in such an eventuality. In fact we feel that the use of the name *Saigonodon* is to be preferred since, unlike *Typhlina*, it has been used several times since its original description (Bocourt 1882 *in* Duméril, Bocourt & Mocquard, 1870-1909: 507; Van Denburgh, 1897: 150 & 1912: 153; Stejneger & Barbour (1917: 73 and 1923: 79); Schmidt, 1922: 682; Klauber, 1931: 336; and Tanner, 1935: 267).
  - 9. The Commission is therefore requested
    - to use its plenary powers to suppress the generic name Typhlina Wagler, 1830 for the purposes of the Law of Priority but not for those of the Law of Homonymy;

(2) to place the generic names

(a) Leptotyphlops Fitzinger, 1843 (gender: masculine), type species by original designation, Typhlops nigricans Schlegel, 1839, and

(b) Ramphotyphlops Fitzinger, 1843 (gender: masculine), type species by original designation, Typhlops multilineatus

Schlegel, 1839,

on the Official List of Generic Names in Zoology:

(3) to place the specific names

- (a) nigricans Schlegel, 1839, as published in the binomen Typhlops nigricans (specific name of type-species of Leptotyphlops Fitzinger, 1843), and
- (b) multilineatus Schlegel, 1839, as published in the binomen Typhlops multilineatus (specific name of type-species of Ramphotyphlops Fitzinger, 1843)

on the Official List of Specific Names in Zoology;

(4) to place the generic name *Typhlina* Wagler, 1830, as suppressed under the plenary powers in (1) above, on the Official Index of Rejected and Invalid Generic Names in Zoology.

## REFERENCES

BOIE, F., 1827. Bemerkungen uber Merrem's Versuch eines Systems der Amphibien. Ite Lieferung: Ophidier. *Isis*, *Jena* 20: 508-566

BOULENGER, G.A., 1890. The Fauna of British India, including Ceylon and Burma. Reptilia and Batrachia. London. xviii + 541 pp.

1892. Zool. Rec. 28 (Rept. Amph.): 1-24

DITMARS, R.L., 1939. A field book of North American snakes. New York. xii + 305 pp. DUMÉRIL, A., BOCOURT, F. & MOCQUARD, F., 1870-1909. Mission scientifique au Mexique et dans l'Amérique centrale. Recherches Zoologiques. Part 3. Sect. 1. Etudes sur les Reptiles. Paris. 1012 pp.

FITZINGER, L., 1826. Neue Classification der Reptilien nach ihren natürlichen Verwandschaften. Vienna, 66 pp.

1843. Systema Reptitium. Vienna. 106 + vi pp.

FITZSIMONS, V.F.M., 1962. Snakes of southern Africa. London. 423 pp.

GOIN, C.J. & GOIN, O.B., 1971. Introduction to Herpetology. 2nd edition. San Francisco. xi + 353 pp.

GRAY, J.E., 1845. Catalogue of the specimens of lizards in the collection of the British Museum. London. 289 pp.

KLAUBER, L.M., 1931. Notes on the worm snake of the Southwest, with descriptions of two new subspecies. *Trans. S. Diego Soc. nat. Hist.* 6: 333-352

LATREILLE, P.A., 1810. Considérations générales sur l'ordre naturel des animaux composant les classes des Crustacés, des Arachnides, et des Insectes; avec un tableau méthodique de leurs genres, disposés en familles. Paris. 444 pp.

LEVITON, A.E., 1971. Reptiles and amphibians of North America. New York. 250 pp. LOVERIDGE, A., 1957. Check list of the reptiles and amphibians of East Africa (Uganda;

Kenya; Tanganyika; Zanzibar). Bull. Mus. comp. Zool. Harv. 117: 153-362

McDOWELL, S.B., 1974. A catalogue of the snakes of New Guinea and the Solomons, with special reference to those in the Bernice P. Bishop Museum, Part 1. Scolecophidia. *J. Herpetol.* 8: 1-57

- MERTENS, R., 1971. Die Herpetofauna sudwest-Afrikas. Abh. senckenb. naturlorsch. Ges. 529: 1-110
- OPPEL, M., 1811. Die Ordnung, Familien und Gattungen der Reptilien als Prodrom einer Naturgeschichte derselben. Munich. xii + 86 pp.
- PARKER, H.W., 1949. The snakes of Somaliland and the Sokotra Islands. Zool. Verh. Leiden 6: 1-115
- PETERS, J.A. & OREJAS MIRANDA, B.R., 1970. Catalogue of the neotropical Squamata: Part 1. Snakes. Bull. U.S. natn. Mus. 297 1: 1-347
- PETERS, W., 1881. Uebersicht der zu den Familien der Typhlopes und Stenostomi gehörigen Gattungen oder Untergattungen. Sber. Ges. naturl. Freunde Berl. 1881: 69-71
- PITMAN, C.R.S., 1974. A guide to the snakes of Uganda. Revised edition. Codicote. 290 pp. PORTER, K.R., 1972. Herpetology. Philadelphia, London and Toronto. xi + 524 pp.
- ROBB, J., 1966. The generic status of the Australasian Typhlopids (Reptilia: Squamata).

  Ann. Mag. nat. Hist. 9: 657-679
- ROZE, J.A., 1966. La Taxonomia y Zoogeografia de los Ofidios en Venezuela. Caracas. 362 pp. SCHLEGEL, H., 1827. Erpetologische Nachrichten. Isis (Jena), 20: 281-294
- ————— 1837-1844. Abbildungen neuer oder unvollständig bekannter Amphibien, nach der Natur oder dem Leben entworlen. Düsseldorf. xiv + 141 pp. [pp. 1-32, 1837; pp. 33-80, 1839; pp. 81-141, 1844]
- SCHMIDT, K.P., 1922. Scientific results of the expedition to the Gulf of California in charge of C.H. Townsend, by the U.S. fisheries steamship 'Albatross', in 1911. Commander G.H. Burrage. U.S.N., commanding VIII. The amphibians and reptiles of Lower California and neighboring islands. Bull. Am. Mus. nat. Hist. 46: 607-707
  - —————— 1953. A check list of North American amphibians and reptiles. 6th edition. Chicago. 280 pp.
- SCHNEJDER, J.G., 1801. Historiae Amphibiorum naturalis et literariae. Jena. Vol. 2. vi + 364 pp.
- SMITH, H.M. & TAYLOR, E.H., 1945. An annotated checklist and key to the snakes of Mexico. Bull. U.S. natn. Mus. 187: 1-239
- SPIX, J. de. 1824. Serpentum brasiliensium species novae ou Histoire naturelle des espèces nouvelles de serpens, recueillies et observées pendant le voyage dans l'intérieur du Brésil dans les années 1817, 1818, 1819, 1820, Exécuté par Ordre de sa Majesté le Roi de Bavière. Monachii. 75 pp.
- STEJNEGER, L., 1891. Notes on some North American snakes. *Proc. U.S. natn. Mus.* 14: 501-505
- reptiles. 2nd edition. London. x + 171 pp.
- TANNER, V.M., 1935. The worm snake, Saigonodon humilis (Baird & Girard) found in Utah. Proc. Utah. Acad. Sci. 12: 267-270
- VAN DENBURGH, J., 1897. The reptiles of the Pacific coast and Great Basin. Occ. Pap. Calif. Acad. Sci. 5: 1-236
- WAGLER, J., 1830. Natürliches System der Amphibien mit vorangehender Classification der Säugethiere und Vögel. Ein Beitrag zur vergleichenden Zoologie. Munich, Stuttgart and Tübingen, vi + 354 pp.
- WITTE, G.F. de, 1953. Reptiles. Explor. Parc natn. Upemba Miss G.F. de Witte 6: 1-322 WRIGHT, A.H. & WRIGHT, A.A., 1957. Handbook of the snakes of the United States and Canada. London. 2 vols. 1105 pp.